

RH  
01  
A1566  
ishes

The Red Alga *Polysiphonia*  
(Rhodomelaceae) in the  
Northern Gulf of California

GEORGE J. HOLLENBERG  
and  
JAMES N. NORRIS

SMITHSONIAN CONTRIBUTIONS TO THE MARINE SCIENCES • NUMBER 1

## SERIES PUBLICATIONS OF THE SMITHSONIAN INSTITUTION

Emphasis upon publication as a means of "diffusing knowledge" was expressed by the first Secretary of the Smithsonian. In his formal plan for the Institution, Joseph Henry outlined a program that included the following statement: "It is proposed to publish a series of reports, giving an account of the new discoveries in science, and of the changes made from year to year in all branches of knowledge." This theme of basic research has been adhered to through the years by thousands of titles issued in series publications under the Smithsonian imprint, commencing with *Smithsonian Contributions to Knowledge* in 1848 and continuing with the following active series:

*Smithsonian Contributions to Anthropology*  
*Smithsonian Contributions to Astrophysics*  
*Smithsonian Contributions to Botany*  
*Smithsonian Contributions to the Earth Sciences*  
*Smithsonian Contributions to the Marine Sciences*  
*Smithsonian Contributions to Paleobiology*  
*Smithsonian Contributions to Zoology*  
*Smithsonian Studies in Air and Space*  
*Smithsonian Studies in History and Technology*

In these series, the Institution publishes small papers and full-scale monographs that report the research and collections of its various museums and bureaux or of professional colleagues in the world of science and scholarship. The publications are distributed by mailing lists to libraries, universities, and similar institutions throughout the world.

Papers or monographs submitted for series publication are received by the Smithsonian Institution Press, subject to its own review for format and style, only through departments of the various Smithsonian museums or bureaux, where the manuscripts are given substantive review. Press requirements for manuscript and art preparation are outlined on the inside back cover.

S. Dillon Ripley  
Secretary  
Smithsonian Institution

The Red Alga *Polysiphonia*  
(Rhodomelaceae) in the  
Northern Gulf of California

*George J. Hollenberg*  
and *James N. Norris*

ISSUED

1 1 1 1  
AUG 25 1977



SMITHSONIAN INSTITUTION PRESS

City of Washington

1977

## ABSTRACT

Hollenberg, George J., and James N. Norris. The Red Alga *Polysiphonia* (Rhodomelaceae) in the Northern Gulf of California. *Smithsonian Contributions to the Marine Sciences*, number 1, 21 pages, 10 figures, 1977.—Taxonomic studies of *Polysiphonia* show 14 species to be present in the northern Gulf of California, Mexico. One of these, *P. sphaerocarpa* var. *cheloniae* is described herein as a new variety, and is found growing exclusively on the green sea turtle, *Chelonia mydas* L. This and three other species are recorded for the first time in the Gulf of California. The distribution of six other species is extended.

OFFICIAL PUBLICATION DATE is handstamped in a limited number of initial copies and is recorded in the Institution's annual report, *Smithsonian Year*. SERIES COVER DESIGN: Seascape along the Atlantic coast.

---

### Library of Congress Cataloging in Publication Data

Hollenberg, George J.

The red alga *Polysiphonia* (Rhodomelaceae) in the northern Gulf of California.

(Smithsonian contributions to the marine sciences ; no. 1)

Bibliography: p.

Supt. of Docs. no.: SI 1.29:37

I. *Polysiphonia*. 2. Algae—Mexico—California, Gulf of. I. Norris, James N., joint author.

II. Title. III. Series: Smithsonian Institution. Smithsonian contributions to the marine sciences ; no. 1.

QK569.R37H64 589'.41 76-608299

## Contents

	<i>Page</i>
Introduction .....	1
Key to the Species of <i>Polysiphonia</i> .....	2
<i>Polysiphonia confusa</i> Hollenberg .....	2
<i>Polysiphonia flaccidissima</i> Hollenberg .....	4
<i>Polysiphonia johnstonii</i> Setchell & Gardner var. <i>johnstonii</i> .....	4
<i>Polysiphonia johnstonii</i> var. <i>concinna</i> (Hollenberg) Hollenberg .....	7
<i>Polysiphonia masonii</i> Setchell & Gardner .....	8
<i>Polysiphonia mollis</i> J. D. Hooker & Harvey in Harvey .....	9
<i>Polysiphonia pacifica</i> var. <i>delicatula</i> Hollenberg .....	10
<i>Polysiphonia paniculata</i> Montagne .....	11
<i>Polysiphonia savatieri</i> Hariot .....	12
<i>Polysiphonia scopulorum</i> var. <i>villum</i> (J. G. Agardh) Hollenberg .....	14
<i>Polysiphonia simplex</i> Hollenberg .....	14
<i>Polysiphonia sonorensis</i> Hollenberg .....	16
<i>Polysiphonia sphaerocarpa</i> var. <i>cheloniae</i> , new variety .....	16
<i>Polysiphonia</i> species .....	17
Literature Cited .....	19
Index .....	21



# The Red Alga *Polysiphonia* (Rhodomelaceae) in the Northern Gulf of California

*George J. Hollenberg  
and James N. Norris*

## Introduction

The earliest records of *Polysiphonia* Greville (1824:308) in the Gulf of California were provided by Setchell and Gardner (1924), who described three new species. Later, Dawson (1944), in his investigations on marine algae of the Gulf of California, recorded seven species, including a new one. A detailed study of the taxa of *Polysiphonia* from the Gulf of California and Pacific coast of Baja California was first presented by Hollenberg (1961). These three papers constitute the basis for our knowledge of *Polysiphonia* in the Gulf of California.

Members of the genus *Polysiphonia* (Rhodomelaceae, Ceramiales) are distinguished by several important taxonomic characters: the number of pericentral cells; presence or absence of cortication; the manner of origin of rhizoids; the manner of origin of branches; the nature and position of trichoblasts; and the nature of spermatangial branchlets. These features have been pointed out in previous investigations by Hollenberg (1942a; 1942b; 1944; 1961; 1968a).

Herein we present our studies of recent intertidal and subtidal collections of *Polysiphonia* from the

northern Gulf of California. These collections were obtained by the junior author during field expeditions in 1972–1974 on the Gulf coasts of Baja California and Sonora. In addition, the following collections were utilized during the course of this study: those of the senior author from Puerto Peñasco, Sonora, in 1967; some additional material of the late Dr. E. Yale Dawson; and those of the junior author and Katina Bucher from the cruise of the R/V *Dolphin* to Las Islas de la Cintura (Midriff Islands) (Norris and Bucher, 1976).

Collectors are designated as follows: JN for J. N. Norris; H for G. J. Hollenberg; D for E. Y. Dawson; and KB for Katina E. Bucher. The numbers cited correspond to the collector's field notebooks. Specimens are being deposited in the following herbaria: first set, U.S. National Herbarium, Smithsonian Institution (US); duplicate material to University of California, Berkeley (UC); Phycology Herbarium, University of Arizona (ARIZ); Universidad Nacional Autónoma de México (MEXU). Herbaria housing the holotypes are abbreviated in accordance with Holmgren and Keuken (1974). Geographical locales under "Specimens Studied" are listed from north to south.

ACKNOWLEDGMENTS.—The Latin diagnosis was kindly provided by Dr. Hannah T. Croasdale (Dartmouth College). Field studies for collections were made possible through the support of the Na-

---

*George J. Hollenberg, Department of Biology, University of Redlands, Redlands, California 92373. James N. Norris, Department of Botany, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560.*

tional Science Foundation (research grant GB-28623) to Dr. M. Neushul and J. Norris (Marine Science Institute, University of California, Santa Barbara). The junior author also wishes to thank again the University of Arizona for use of their Laboratorio de Biología Marina, Puerto Peñasco, Sonora, while he served as resident Marine Biologist/Station Director of the facility. Collections by Norris and Bucher from Las Islas de la Cintura were made possible only through the generosity of Dr. William Fenical (Scripps Institution of Oceanography), Chief Scientist on the R/V *Dolphin* cruise of April 1974.

Collections from the sea turtles were obtained by Dr. Richard S. Felger (Arizona-Sonora Desert Museum, Tucson, Arizona) and Edward W. Moser (Summer Institute of Linguistics, University of North Dakota). Further collections were given to Norris by Dr. Felger, Professor Richard Evans Schultes (Director, Botanical Museum of Harvard University), and Dr. Andrew T. Weil (Botanical Museum of Harvard University). Finally, we wish to thank Drs. Isabella A. Abbott, Robert W. Hoshaw, and Katina Bucher for reading the manuscript and offering suggestions for its improvement.

### Key to the Species of *Polysiphonia*

THE SPECIES ARE ARRANGED ALPHABETICALLY IN THE TEXT

1. Pericentral cells 4 (Section Oligosiphonia) ..... 2  
Pericentral cells more than 4 (Section Polysiphonia) ..... 11
2. Trichoblasts or scar-cells in general, regularly 1 per segment ..... 3  
Trichoblasts or scar-cells lacking or not regularly 1 per segment ..... 9
3. Branching dichotomous throughout ..... *P. masonii*  
Branching lateral, not strictly dichotomous ..... 4
4. Growing on sea turtles ..... *P. sphaerocarpa* var. *cheloniae*, new variety  
Not growing on sea turtles ..... 5
5. Branches arising from a primary branch of a trichoblast (axillary to a trichoblast) .....  
..... *P. flaccidissima*  
Branches arising from an entire trichoblast primordium ..... 6
6. Epiphytic, mostly less than 2 cm high ..... *P. savatieri*  
Usually not epiphytic ..... 7
7. Segments in main branches mostly shorter than their diameter or sometimes as long .....  
..... *P. simplex*  
Segments in main branches mostly 1.5 times as long as their diameter or longer ..... 8
8. Trichoblasts usually well developed; spermatangial stichidia arising as a primary branch of a trichoblast ..... *P. mollis*  
Trichoblasts very rudimentary; spermatangial stichidia arising from an entire trichoblast primordium ..... *Polysiphonia* species
9. Thalli delicate, mostly intertidal, commonly forming mats or furry patches .....  
..... *P. scopulorum* var. *villum* ..... 10  
Thalli not as delicate, mostly subtidal, not forming mats or patches
10. Thalli mostly less than 2 cm high; trichoblasts infrequent to rare ..... *P. pacifica* var. *delicatula*  
Thalli up to 4 cm high, of a terracotta color; trichoblasts numerous, arising at irregular intervals ..... *P. sonorensis*
11. Pericentral cells 5-6 ..... 12  
Pericentral cells 8 or more ..... 13
12. Thalli occasionally epiphytic, mostly 5 cm or more high ..... *P. johnstonii* var. *johnstonii*  
Thalli usually epiphytic, mostly less than 5 cm high ..... *P. johnstonii* var. *concinna*
13. Thalli mostly less than 2 cm high ..... *P. confusa*  
Thalli mostly much larger, to 20 cm high ..... *P. paniculata*

### *Polysiphonia confusa* Hollenberg

FIGURE 1

*Polysiphonia confusa* Hollenberg, 1961:350, pl. 1; fig. 5.—Dawson, Acelto, and Foldvik, 1964:87, pl. 33d.—Abbott and Hollenberg, 1976:696, fig. 643.

*Polysiphonia inconspicua* sensu Hollenberg, 1944:479 [not *Polysiphonia inconspicua* Reinsch, 1888:146].

DESCRIPTION.—Thalli 0.8–1.5(–3.0) cm high, with erect axes mostly assurgent from prostrate branches 100–175  $\mu$ m in diameter, composed of segments






FIGURE 1.—*Polysiphonia confusa*: A, upper portion of thallus (JN-5577); B, cystocarpic plant (JN-3835).

about as long as the diameter or somewhat longer, attached by numerous unicellular rhizoids, arising 1 to several per segment as cells cut off from the proximal ends of the pericentral cells, and frequently having digitate apices; erect branches 60–150  $\mu\text{m}$  in diameter, the segments 1.0–2.5 times as long; lateral branches arising exogenously in connection with trichoblasts at irregular intervals; pericentral cells 8–10, without cortication; trichoblasts to 1 mm long, with 1 or 2 dichotomies, arising 1 per segment in spiral sequence with  $\frac{1}{4}$  divergence, soon deciduous, leaving persistent scar-cells.

Tetrasporangia, 60–80  $\mu\text{m}$  in diameter, in long spiral series in slightly swollen segments of ultimate and subultimate branches; cystocarps globular, to 225  $\mu\text{m}$  in diameter; spermatangial plants unknown.

TYPE-LOCALITY.—Mid-intertidal rocks, Corona del Mar, Orange County, California.

HOLOTYPE.—H-3285, 24 October 1942, in herbarium of G. J. Hollenberg (now at US).

GULF OF CALIFORNIA DISTRIBUTION.—Mid-intertidal regions; epizoid on sand dollars, epiphytic on *Codium* and growing on hard substrate of buoy; Puerto Peñasco to Desemboque de San Ignacio.

PACIFIC COAST DISTRIBUTION.—U.S.A.: southern California. Mexico: northern Baja California.

SPECIMENS STUDIED.—SONORA: JN-5577, sterile, growing on a sand dollar, *Encope grandis*, Bahía Cholla near Punta Pelicano, legit J. N. and K. B., 25 March 1974; JN-3835, cystocarpic, on NASA buoy, anchored off Laboratorio de Biología Marina, Playa Estación, Puerto Peñasco, legit J. N., 22 March 1973; JN-5158, growing on *Codium* sp., Desemboque de San Ignacio, legit R. E. Schultes, R. S. Felger, and A. T. Weil, 3 April 1974.

REMARKS.—Previously known on the Pacific coast from southern California and northern Baja Cali-

fornia (Hollenberg, 1961:351), this taxon is now recorded in the Gulf of California. It is apparently rare in the northern Gulf; only three collections have come to our attention. Although none of these specimens is from the habitat described as characteristic of *P. confusa* Hollenberg, size and structural features leave little room for doubt as to the identity as given.

### *Polysiphonia flaccidissima* Hollenberg

FIGURE 2E

*Polysiphonia flaccidissima* Hollenberg, 1942b:783, figs. 8, 19; 1961:351, pl. 2: fig. 2; 1968a:63, figs. 2A, 11.—Dawson, Aceldo, and Foldvik, 1964:88, pl. 78b.—Abbott and Hollenberg, 1976:688, fig. 634.

DESCRIPTION.—Epiphytic or saxicolous algae 10–25 mm high, erect from creeping basal branches attached by numerous unicellular rhizoids, which are cut off as separate cells from the pericentral cells; main erect axes 70–80  $\mu$ m in diameter, with segments mostly 1–2 times as long as the diameter; branches of several to many orders, arising exogenously from a primary branch of a trichoblast, mostly at intervals of 6 segments or less, narrowed gradually below and above, at first usually strict and curving toward the parent branch; pericentral cells 4, totally ecorticate, with thin hyaline walls; trichoblasts simple or mostly with a single dichotomy, tapering to slender apices, arising 1 per segment in spiral sequence, with  $\frac{1}{4}$  divergence, soon deciduous, leaving persistent scar-cells.

Tetrasporangia 50–70  $\mu$ m in diameter, in spiral series in smaller branches, somewhat distending the segments; cystocarps globular; spermatangial stichidia arising as a primary branch of a trichoblast, with 1–2 celled sterile apices.

TYPE-LOCALITY.—On a coralline alga, rocky point, Laguna Beach, Orange County, California.

HOLOTYPE.—H-2269, 1 January 1938, in herbarium of G. J. Hollenberg (now at US).

GULF OF CALIFORNIA DISTRIBUTION.—Intertidal to subtidal, on rocks or occasionally epiphytic; Guaymas to Isla Espiritu Santo.

PACIFIC COAST DISTRIBUTION.—U.S.A.: southern California; Hawaiian Islands. Mexico: northern Baja California; Revilla Gigedo Archipelago; Guerrero. Costa Rica. Panama. Peru.

SPECIMENS STUDIED.—SONORA: JN-4267, tetrasporangial, and JN-4270, on *Padina*, from 1.5–6.1 m depth, Nueva Guaymas, legit Dale Evensen, 13 Oct 1973.

REMARKS.—In the northern Gulf this species is apparently rare, restricted in distribution to Guaymas and vicinity, Sonora. Across the Gulf it has been recorded from Santa Rosalía, and in the southern Gulf from Puerto Escondido and Isla Espiritu Santo (Hollenberg, 1961:352).

### *Polysiphonia johnstonii* Setchell & Gardner var. *johnstonii*

FIGURES 2B, 3

*Polysiphonia johnstonii* Setchell & Gardner, 1924:767.—Hollenberg, 1944:475; 1961:357, pl. 2: figs. 4, 5.—Dawson, 1944:329; 1949:237; 1951:53, 57; 1954b:160; 1959:32; 1966:29.—Dawson, Neushul, and Wildman, 1960a:72; 1960b:26.—Norris, 1972:16.—Abbott and Hollenberg, 1976:699, fig. 649.

*Polysiphonia marchantae* Setchell & Gardner, 1924:768, pl. 49a.

*Polysiphonia sinicola* Setchell & Gardner, 1924:769.

*Polysiphonia acuminata* sensu Dawson, 1950:68; 1954b:159 [non *Polysiphonia acuminata* Gardner, 1927:100].

*Polysiphonia richardsoni* sensu Segi, 1951:253.—Dawson, 1954b:160 [in part, with reference to Gulf of California specimens cited by Setchell and Gardner, 1924:768 as *P. marchantae* only; non *Polysiphonia richardsoni* W. J. Hooker ex Harvey in W. J. Hooker, 1833:333].

DESCRIPTION.—Thalli relatively rigid, 5–12 cm high, dull red in color, nearly black on drying, attached by numerous unicellular rhizoids, which are cut off as separate cells from the pericentral cells of a limited prostrate base; primary erect branches several, relatively distinct, to 1 mm in diameter, repeatedly branched, the branches spirally arranged, arising at a narrow angle, which widens as the branch matures; ultimate branches usually relatively close and dense; segments in main branches 1.5 times as long as the diameter or shorter, with 5–6 pericentral cells, totally ecorticate; trichoblasts once or twice forked, arising 1 per segment in spiral sequence, with  $\frac{1}{5}$  or  $\frac{1}{6}$  divergence, soon deciduous, leaving persistent scar-cells; branches replacing trichoblasts in origin at irregular intervals.

Tetrasporangia 1(–2) per segment, 70–90  $\mu$ m in diameter; cystocarps globular, 450–500  $\mu$ m in diam-




FIGURE 2.—A, *Polysiphonia johnstonii* var. *concinna*, branch tip of a tetrasporic plant (after Hollenberg, 1944); B, *P. johnstonii* var. *johnstonii*, apical portion of an axis showing trichoblasts and characteristically incurved tips (after Hollenberg, 1961); C, *P. sonorensis*, branch tip of the holotype (after Hollenberg, 1961); D, *P. masonii*, upper portion of a tetrasporic plant (after Hollenberg, 1961); E, *P. flaccidissima*, branch tip (after Hollenberg, 1942b).




FIGURE 3.—*Polysiphonia johnstonii* var. *johnstonii*: A, B, habits of two specimens showing the divergent manner of branching (JN-3201).

eter, nearly sessile; spermatangial stichidia arising as a primary branch of a trichoblast, 100–140 by 30–40  $\mu\text{m}$  in length and diameter, respectively.

**TYPE-LOCALITY.**—On *Gracilaria* sp., Isla San Esteban, Gulf of California, Mexico.

**HOLOTYPE.**—Ivan M. Johnston 110, April 1921 (CAS 1361, now at UC).

**GULF OF CALIFORNIA DISTRIBUTION.**—Intertidal and subtidal, on rocks or epiphytic; Puerto Peñasco to La Paz.

**PACIFIC COAST DISTRIBUTION.**—Santa Catalina Island, California; northern Baja California; Sinaloa.

**SPECIMENS STUDIED.**—SONORA: JN-3102, rocky intertidal, Punta Pelicano, 30 Jun 1972; JN-3260, on *Sargassum* sp. on rock platform with sand areas, Playa Hermosa, Puerto Peñasco, legit J. N. and K. B., 26 Jul 1972; JN-4040, spermatangial, intertidal rock platform, Playa Hermosa, Puerto Peñasco, legit J. N. and K. B., 3 May 1973; JN-5150, beach drift after strong winds, Desemboque de San Ignacio, legit R. E. Schultes, R. S. Felger, and A. T. Weil, 3 Apr 1974. BAJA CALIFORNIA DEL NORTE: JN-5462, 1–7 m depth, Punta la Gringa, Bahía de Los Angeles, legit J. N. and K. B., 28 Apr 1974; JN-3029, 5 m depth, Punta la Gringa, Bahía de los Angeles, 22 May 1972; JN-3201, and JN-3209 (a slender form), estuary, Bahía de las Animas, legit J. N. and G. Boehlert, 23 May 1972. LAS ISLAS DE LA CINTURA: JN-5280, tetrasporangial and cystocarpic, 10 m depth, W side of Roca Blanca, Puerto Refugio, Isla Ángel de la Guarda, legit J. N., J. Paul, and K. Robertson, 21 Apr 1974; JN-5314, intertidal to 10 m depth, Puerto Refugio, Isla Ángel de la Guarda, legit J. N. and K. B., 21 Apr 1974; JN-5369, 5–12 m depth off small islet, Puerto Refugio, Isla Ángel de la Guarda, 21 Apr 1974; JN-5762, intertidal, rocky shore NE side of Puerto Refugio, Isla Ángel de la Guarda, 23 Apr 1974; JN-5483, 3–10 m depth, S end of Isla Estanque, legit J. N. and K. B., 27 Apr 1974; JN-5508 and JN-5616A, 3–12 m depth, S end of Isla Estanque, legit J. N. and K. B., 27 Apr 1974; JN-5561, tetrasporangial, 3–8 m depth, rocky shore SE end of Isla San Esteban, 25 Apr 1974; JN-5528, cystocarpic, 3–5 m depth, off SE end of Isla San Esteban, 26 Apr 1974; JN-5707, epiphytic on *Laurencia sinicola*, and JN-5709, 3–8 m depth, SE end of Isla San Esteban, 25 Apr 1974; JN-5518, 1–5 m depth, SE end of Isla San Esteban, legit K. B., 26 Apr 1974.

**REMARKS.**—One of the most frequently encountered species of *Polysiphonia* in the northern Gulf, its distribution is now extended northward to Puerto Peñasco from Puerto Libertad (Hollenberg, 1961:358). In the northern Gulf, specimens are usually epiphytic and commonly sterile.

*Polysiphonia johnstonii* var. *concinna*  
(Hollenberg) Hollenberg

FIGURES 2A, 4D

*Polysiphonia concinna* Hollenberg, 1944:474, fig. 10.—Dawson, 1951:53, 56; 1954b:159; 1959:32.

*Polysiphonia johnstonii* var. *concinna* (Hollenberg) Hollenberg, 1961:358, pl. 3: fig. 4.—Abbott and Hollenberg, 1976: 699, fig. 650.

**DESCRIPTION.**—This variety is similar to *P. johnstonii* var. *johnstonii* in detailed features, but smaller and shorter in dimensions throughout.

**TYPE-LOCALITY.**—Epiphytic on other algae, near Scripps Institution of Oceanography, La Jolla, San Diego County, California.

**HOLOTYPE.**—H-2015, December 1936, in herbarium of G. J. Hollenberg (now at US).

**GULF OF CALIFORNIA DISTRIBUTION.**—Mid-intertidal to subtidal, epiphytic on various algae, occasionally on rocks; Puerto Peñasco to Bahía Agua Verde.

**PACIFIC COAST DISTRIBUTION.**—Southern California; northern Baja California; Sinaloa.

**SPECIMENS STUDIED.**—SONORA: H 67–64, spermatangial, cystocarpic, on *Sargassum* sp. in drift, Playa Tucson, Punta Pelicano, near Puerto Peñasco, 8 Apr 1967; H 67–70, on *Sargassum* sp. in drift, Playa Estación, Puerto Peñasco, 8 Apr 1967. BAJA CALIFORNIA DEL NORTE: JN-4667, and JN-4668 spermatangial, and JN-4669 cystocarpic, the latter two on *Euclidean uncinate*, and JN-4670, cystocarpic, on *Codium* sp., 3–5 m depth N side of Puerto Calamajue, legit J. N. and K. B., 28 Mar 1973; JN-3011, cystocarpic, Islas de los Gemelos, Bahía de los Angeles, 21 May 1972; JN-4752, on *Padina*, Isla de los Gemelos, Bahía de los Angeles, 22 Dec 1972; JN-3238, spermatangial, on *Codium* sp., Bahía San Francisquito, 24 May 1972. LAS ISLAS DE LA CINTURA: JN-5804, cystocarpic, 3–6 m depth, off W side of Roca Blanca, Puerto Refugio, Isla Ángel de la Guarda, 22 Apr 1974; JN-5616B, 1–10 m depth off S end of Isla Estanque at the S end of Isla Ángel de la Guarda, legit J. N. and K. B., 27 Apr 1974; JN-5520b, on *Codium* sp., 1–5 m depth, SE end, off Isla San Esteban, legit K. B., 26 Apr 1974.

**REMARKS.**—Previously encountered off Isla San Jorge in the upper Gulf (Hollenberg, 1961:358), it was not surprising to find *P. johnstonii* var. *concinna* in Puerto Peñasco and vicinity. Number JN-3011, cited above, is placed in this variety with some hesitation, since the branches frequently arise in pairs one segment apart. Also the procarpal branches are numerous and present an unusual appearance. They seemingly arise from relatively




FIGURE 4.—A-C, *Polysiphonia sphaerocarpa* var. *cheloniae* (JN-4765): A, the upper branching and a young cystocarp; B, a nearly mature cystocarp; C, two spermatangial stichidia and the related trichoblast of one. D, *P. johnstonii* var. *concinna*, several young cystocarps with seemingly arrested development (JN-3011).

large trichoblast primordia (scar-cells) and remain prominent and undeveloped, at least for the most part.

### *Polysiphonia masonii* Setchell & Gardner

#### FIGURE 2D

*Polysiphonia masonii* Setchell & Gardner, 1930:160.—Hollenberg, 1942b:783; 1961:358, pl. 6: figs. 2, 3.—Dawson, 1954b:160.

DESCRIPTION.—Reddish brown epiphytes 2–3 cm high, flaccid, attached by a bundle of short unicellular rhizoids with digitate apices, arising from

the lower segments of the erect primary axis; main axis 350–400 μm in diameter below, repeatedly and regularly branched dichotomously, composed of segments mostly as long as the diameter; branches nearly as large as the main axis at the point of insertion, arising from an entire trichoblast primordium; pericentral cells 4, ecorticate; trichoblasts to 480 μm long, with 2 or 3 dichotomies, tapering to very slender apices, arising 1 per segment in spiral sequence with  $\frac{1}{4}$  divergence, soon deciduous, leaving persistent scar-cells.

Tetrasporangia in spiral series in the ultimate branches, 85–95 μm in diameter, not much distending the segments; cystocarps subspherical nearly

sessile, 300–325  $\mu\text{m}$  in diameter; spermatangial stichidia arising from a primary branch of a trichoblast, broadly fusiform.

TYPE-LOCALITY.—Epiphytic on *Zostera*, Isla Guadalupe, off the Pacific Coast of Baja California.

HOLOTYPE.—H. L. Mason 86, April 1925 (CAS 173618, now at UC).

GULF OF CALIFORNIA DISTRIBUTION.—Intertidal, Puerto Peñasco.

PACIFIC COAST DISTRIBUTION.—Isla Guadalupe.

SPECIMEN STUDIED.—SONORA: H 67–60, tetrasporangial, intertidal, Playa Estación, Puerto Peñasco, 8 Apr 1967.

REMARKS.—Now reported for the first time in the Gulf of California, this seems to be the only record of the species other than those recorded from Isla Guadalupe (Hollenberg, 1961:359). Rarely encountered, it is known only from the present Gulf collection; three others, including the type specimen, from Isla Guadalupe.

### *Polysiphonia mollis* J. D. Hooker & Harvey in Harvey

FIGURE 5B

*Polysiphonia mollis* J. D. Hooker & Harvey in Harvey, 1847:43.—Dawson, 1957:8; 1966:29.—Dawson, Neushul, and Wildman, 1960a:72.—Hollenberg, 1961:359, pl. 4: fig. 2.—Abbott and Hollenberg, 1976:688, fig. 636.

*Polysiphonia eastwoodae* Setchell & Gardner, 1930:161.—Hollenberg, 1948:160.—Dawson, 1954b:160.

*Polysiphonia tongatensis* var.? sensu Setchell & Gardner, 1930:160 [non *Polysiphonia tongatensis* Harvey in Kuetzing, 1864:14].

*Polysiphonia snyderae* Kylin, 1941:35, pl. 12: fig. 34.—Hollenberg, 1942a:784, fig. 9.—Dawson, 1944:330; 1951:56; 1954b:160.

*Polysiphonia snyderae* var. *heteromorpha* Hollenberg, 1942b:784.

*Polysiphonia snyderae* var. *intricata* Hollenberg, 1942b:785.—Dawson, 1954b:161.

*Polysiphonia tongatensis* sensu Segi, 1951:207 [in part, with reference to Revilla Gigedo Islands' material of Setchell and Gardner, 1930:160 only; non *Polysiphonia tongatensis* Harvey in Kuetzing, 1864:14].

DESCRIPTION.—Thalli dull reddish brown, mostly 5–12 cm high from a discoid base or mostly assurgent from prostrate branches of limited extent, attached by unicellular rhizoids with frequently digitate apices; arising 1–2(–5) per segment from the proximal ends of the pericentral cells, from

which they are cut off as separate cells by a curving wall; erect branches 300–400  $\mu\text{m}$  in diameter below, commonly 60–180  $\mu\text{m}$  in diameter in upper parts; segments in main branches mostly 1–2 times as long as the diameter, but sometimes considerably longer; main erect branches richly branched pseudodichotomously above, mostly naked below; pericentral cell 4, ecorticate; trichoblasts 1 per segment in spiral sequence, with  $\frac{1}{4}$  divergence, soon deciduous, leaving persistent scar-cells; branches arising at irregular intervals, commonly 6–10 segments apart, a branch replacing a trichoblast in the spiral sequence.

Tetrasporangia 60–70  $\mu\text{m}$  in diameter, in spiral series in the ultimate and subultimate branches; cystocarps abundant, ovoid to nearly globular, 300–350  $\mu\text{m}$  in diameter; spermatangial stichidia comprising a primary branch of a trichoblast.

TYPE-LOCALITY.—Epiphytic on other algae, Tasmania.

HOLOTYPE.—Collection of R. Gunn labeled "original sp. . . parasitical on large algae" in Harvey Herbarium (TCD).

GULF OF CALIFORNIA DISTRIBUTION.—Usually intertidal, epiphytic, or growing on rock, wood, shells, etc.; El Tornillal to Bahía San Lucas.

PACIFIC COAST DISTRIBUTION.—Southern British Columbia to northern Baja California; Sinaloa.

SPECIMENS STUDIED.—SONORA: JN-3326, adrift, inlet, El Tornillal, legit D. L. Lindquist; H 67–1.5, legit T. Scalone, Playa Estación, Puerto Peñasco, 28 Jan 1967; D27210 (US), rocky reef, Playa Tucson, N side of Punta Pelicano, vicinity of Puerto Peñasco, legit E. Y. Dawson, 6 Apr 1966. BAJA CALIFORNIA DEL NORTE: JN-2998a, spermatangial, from 7.6 m depth, Isla la Ventana, Bahía de los Angeles, 21 May 1972; JN-3265a, growing on "pen shell," *Pinna rugosa*, 3–4.5 m depth, Bahía San Francisquito, 24 May 1972; JN-3226, a slender form, growing on shells, Bahía San Francisquito, 24 May 1972.

REMARKS.—The recent collection from El Tornillal represents a slight northward extension from Puerto Peñasco, Sonora (Dawson, 1966:29). This species is widely distributed in temperate and subtropical regions. It is most frequently encountered in sheltered waters.

In collection JN-3265a the segments in the main axes are shorter than the typical length for the species. A more unusual feature in JN-2998a is the fact that the spermatangial stichidia arise from the entire trichoblast primordium. In both JN-2998a




FIGURE 5.—A, *Polysiphonia paniculata*, branch tip (after Hollenberg, 1961); B, *Polysiphonia mollis*, branch tip (after Hollenberg, 1961).

and JN-3265a the trichoblasts are very rudimentary or are represented by scar-cells only. Furthermore, rhizoids in JN-3265a are cut off as separate cells from the pericentral cells. The latter feature does not normally occur in species such as *P. pacifica* in which spermatangial stichidia arise from the entire trichoblast primordium. More material representing these two entities is needed for confident identification.

***Polysiphonia pacifica* var. *delicatula* Hollenberg**

FIGURE 6B

*Polysiphonia pacifica* var. *delicatula* Hollenberg, 1942b:778, 1961:362.—Smith, 1944:360.—Abbott and Hollenberg, 1976: 690.

DESCRIPTION.—Usually medium red, 0.5–2.0 cm high from prostrate branches 50–110  $\mu\text{m}$  in diameter, arising endogenously from prostrate branches,






FIGURE 6.—A, *Polysiphonia sphaerocarpa* var. *cheloniae*, branch apices of a spermatangial plant (JN-4765); B, *P. pacifica* var. *delicatula*, branch apices (JN-4848).

and composed of segments 1.0–2.0 (–3.0) times as long; branches short and mostly with blunt apices; pericentral cells 4, ecorticate; trichoblasts lacking or exceedingly rare.

Tetrasporangia 50–60  $\mu\text{m}$  in diameter, in straight series; cystocarps slightly urceolate; spermatangial stichidia slender, 140–200  $\mu\text{m}$  in length by 30–40  $\mu\text{m}$  in diameter, on very short pedicels, leaving persistent scar-cells when shed.

TYPE-LOCALITY.—On wharf piling, low tide level, Municipal Wharf, Monterey, Monterey County, California.

HOLOTYPE.—H 2888, 20 July 1939, in herbarium of G. J. Hollenberg (now at US).

GULF OF CALIFORNIA DISTRIBUTION.—Epiphytic on various algae or growing on rocks, low intertidal to subtidal; Puerto Peñasco to Guaymas; Bahía San Lucas.

PACIFIC COAST DISTRIBUTION.—Central California; Chiapas; Galápagos Archipelago.

SPECIMENS STUDIED.—SONORA: H 67–53, on coralline algae, Playa Estación, Puerto Peñasco, 25 Mar 1967; JN-4848a, epiphytic on *Gelidium johnstonii* (JN-4827), 3.3 m depth, on rock reef, Cabo Lobos, vicinity of Puerto Libertad, legit J. N. and K. B., 17 Nov 1973.

REMARKS.—Seemingly rare in the Gulf of California, its distribution is now extended northward to Puerto Peñasco from Bahía Guaymas (Hollenberg, 1961: 362). This species is generally found on rocks, wood, or epiphytic in deep or quiet waters.

### *Polysiphonia paniculata* Montagne

FIGURES 4A, 5A, 7

*Polysiphonia paniculata* Montagne, 1842:254.—Kuetzing, 1863: 15, pl. 45c-e.—Howe, 1914:142.—Hollenberg, 1944:480, fig. 7; 1961:362, pl. 7: fig. 1.—Dawson, Neushul, and Wildman, 1960b:26.—Dawson, Aceldo and Foldvik, 1964:89, pls. 78A, 79.—Dawson, 1954b:160; 1966:29.—Smith, 1969:115 [725].—Abbott and Hollenberg, 1976:701, fig. 652.

*Polysiphonia californica* Harvey, 1853:48.—Kylin, 1941:36.—Dawson, 1944:332.—Smith, 1944:362, pl. 93: fig. 1.

DESCRIPTION.—Thalli densely tufted, soft, dark brownish red, mostly saxicolous, 10–25 cm high, arising from densely matted prostrate branches, attached by numerous unicellular rhizoids, with usually digitate apices, arising 2 or 3 per segment from the proximal end of the pericentral cells, from which they are cut off as separate cells; erect branches arising endogenously from the prostrate branches at close intervals, loosely branched to relatively naked below, densely clothed with branches above; branches of all orders exogenous, the ultimate branches commonly densely plumose, narrowed at the base; pericentral cells mostly 10–12, but sometimes as many as 14 in older parts or as few as 8 in younger parts, totally ecorticate; main axes 300–430  $\mu\text{m}$  in diameter and composed of segments 2.0–2.5(–4.0) times as long; trichoblasts to 800  $\mu\text{m}$  long, simple or mostly with 1 or 2 dichotomies, arising 1 per segment in spiral sequence with  $\frac{1}{4}$  divergence, tardily deciduous, leaving relatively



FIGURE 7.—*Polysiphonia paniculata*, tetrasporic specimen (JN-4552) from the mud/sand flats of Bahía Cholla (vicinity of Puerto Peñasco), Sonora.

large scar-cells; branches arising in association with trichoblasts.

Tetrasporangia 80–100  $\mu\text{m}$  in diameter, more or less spiraling in the ultimate and subultimate branches; cystocarps globular-ovoid, somewhat truncate, 350–400  $\mu\text{m}$  in diameter; spermatangial stichidia 130–250  $\mu\text{m}$  by 50–70  $\mu\text{m}$  without sterile apex at maturity; thalli adhering to paper when dried.

TYPE-LOCALITY.—Peru.

HOLOTYPE.—Collection by Gaudichaud labeled “in fronde Ulvae nematoideae, ad littora Peruviana” in Montagne Herbarium (PC).

GULF OF CALIFORNIA DISTRIBUTION.—Mid-intertidal to subtidal, usually in protected bays and estuaries, Puerto Peñasco to Guaymas.

PACIFIC COAST DISTRIBUTION.—British Columbia to northern Baja California. Peru.

SPECIMENS STUDIED.—SONORA: D 27254, slender form, mud flats, Bahía Cholla, 6 Apr 1966; JN-4551, cystocarpic, sand flats, 1 m depth, Bahía Cholla, 8 Dec 1973; JN-4552, tetrasporic, growing on living sand dollar (*Encope grandis*), sand flats, Bahía Cholla, legit J. N. and J. Kudenov, 8 Dec 1973; JN-4993, sand flats, Bahía Cholla, legit J. N. and K. B., 9 Mar 1974; JN-3697, and JN-3713, both tetrasporangial, intertidal rock platform, W of Playa Hermosa, Puerto Peñasco, 2 Feb 1973; JN-4718, tetrasporangial, attached to sea shells, midway between Isla Tiburón and W of Campo Viboras, Sonora, legit G. L. Kooyman and E. Senate, 19 Dec 1973; JN-4773.5, spermatangial, collected along with *P. sphaerocarpa* var. *chelonina* (JN-4773), in the Infiernillo Channel between Campo Ona and Viboras, Sonora, legit R. S. Felger and E. Moser from a green sea turtle (♀ *Chelonia mydas*) that was harpooned by Guadalupe Lopez, a Seri Indian, 21 Jan 1974.

REMARKS.—This species apparently occurs only in the northern Gulf. Here it has been found mostly in calm waters of estuaries, mud flats, protected bays, and coastal areas. It is also one of the marine algae known to be epizoid on the green sea turtle, *Chelonia mydas* L. in the Gulf (see Felger, Clifton, and Regal, 1976; Norris, unpublished data).

### *Polysiphonia savatieri* Hariot

FIGURE 8A

*Polysiphonia savatieri* Hariot, 1891:226.—Segi, 1951:202, fig. 10a-n, pl. 3: fig. 8.—Dawson, 1954b:160.—Hollenberg, 1961:363, pl. 7: fig. 4; 1968a:77, figs. 37, 38.—Abbott and Hollenberg, 1976:692, fig. 639.

*Polysiphonia minutissima* Hollenberg, 1942b:781, fig. 21.




FIGURE 8.—A, *Polysiphonia savatieri*, upper portion of a tetrasporic thallus (JN-4195); B, *P. scopulorum* var. *villum*, entire thallus (JN-5098A); note the dichotomous branched trichoblast.

**DESCRIPTION.**—Thalli epiphytic, dull reddish brown, mostly 3–10 mm high from a tuft or cushion of rhizoids, usually penetrating the host; rhizoids up to 1 mm long, cut off by a curving wall from the proximal end of the pericentral cells; main axes indistinct, commonly assurgent from the basal attachment, up to 148  $\mu\text{m}$  in diameter; branches of several orders, exogenous, not associated with trichoblasts in origin, arising at variable intervals; pericentral cells 4, ecorticate; trichoblasts relatively short, 1 per segment in spiral sequence, with  $\frac{1}{4}$  divergence, soon deciduous, leaving persistent scar-cells.

Tetrasporangia in short spiral series, up to 80  $\mu\text{m}$  in diameter, somewhat distending the segments; cystocarps slightly urceolate or mostly globular at maturity, 225–290  $\mu\text{m}$  in diameter; spermatangial stichidia about 135  $\mu\text{m}$  long and 40  $\mu\text{m}$  in diameter,

comprising one primary branch of a trichoblast, without sterile apex.

**TYPE-LOCALITY.**—Epiphytic on large algae, Yokosuka, Japan.

**HOLOTYPE.**—A collection by Dr. Savatier, about 1885 (PC).

**GULF OF CALIFORNIA DISTRIBUTION.**—Mid-intertidal to shallow subtidal, epiphytic on various algae, growing on rocks or in tide pools; Puerto Peñasco to Bahía de Las Animas.

**PACIFIC COAST DISTRIBUTION.**—Central California to northern Baja California; Isla Guadalupe, Japan. Western tropical Pacific.

**SPECIMENS STUDIED.**—SONORA: JN-5030, on *Neogardhiella baileyii*, Punta Pelicano, legit J. N., K. B., and D. Moore, 17 Mar 1974; JN-3182, tetrasporangial, on intertidal rocks, W of Casa García, Playa Estación, Puerto Peñasco, legit J. N. and K. B., 11 Jul 1972; JN-3545, cystocarpic, Cumpleaños tide

pool, Playa Estación, Puerto Peñasco, legit J. N. and K. B., 8 Sep 1972; JN-4537, tetrasporangial, Cumpleaños tide pool, Playa Estación, Puerto Peñasco, legit J. N. and K. B., 9 Nov 1973; JN-4195, tetrasporangial, and JN-4197, cystocarpic, on intertidal platform and tide pools, Playa Estación, Puerto Peñasco, 1 Jun 1973; H 67-14, low intertidal, Playa Estación, Puerto Peñasco, 25 Feb 1967; JN-3878, spermatangial, on *Codium* sp. in front of Laboratorio de Biología Marina, Playa Estación, Puerto Peñasco, 14 Apr 1973; JN-4266, cystocarpic, 6 m depth, Neuva Guaymas, legit D. Evanson, 13 Oct 1973. BAJA CALIFORNIA DEL NORTE: JN-3126, epiphytic on *Sargassum*, 9 m depth, Rocas Consag, legit D. L. Lindquist, 2 Jun 1972; JN-3281, spermatangial, on *Codium* sp., Puertecitos, legit J. N. and G. Boehlert, 17 May 1972; JN-4448, Isla la Ventana, Bahía de Los Angeles, 27 Jul 1973; JN-4442, on *Dictyota* sp., rocky intertidal, Isla la Ventana, Bahía de Los Angeles, 27 Jul 1973; JN-3216b, tetrasporangial, on *Gracilaria* sp., Bahía de Las Animas, legit J. N. and G. Boehlert, 23 May 1972.

REMARKS.—This species is infrequently encountered throughout its Pacific coast distribution (Hollenberg, 1961:364). It may now be added to the northern Gulf of California marine flora, where it was rather frequent among the recent collections.

*Polysiphonia scopulorum* var. *villum*  
(J. G. Agardh) Hollenberg

FIGURE 8B

*Polysiphonia scopulorum* var. *villum* (J. G. Agardh) Hollenberg, 1968a:81, fig. 7A.—Abbott and Hollenberg, 1976:692, figs. 640, 641.

*Polysiphonia villum* J. G. Agardh, 1863:941.

*Lophosiphonia villum* (J. G. Agardh) Setchell & Gardner, 1903:329.—Hollenberg, 1942a:535.—Dawson, 1944:332, pl. 48: figs. 1-6; 1951:53; 1954b:161; 1963:421, pl. 169: figs. 1-3.—Taylor, 1945:304.

*Lophosiphonia scopulorum* sensu Dawson, 1959:36 [non *Lophosiphonia scopulorum* (Harvey) Womersley, 1950:188].

DESCRIPTION.—Brownish red thalli mostly 5–10 mm high from prostrate branches attached by unicellular rhizoids arising from the center of the pericentral cells, from which they are not cut off by cross walls; erect branches unbranched or mostly very sparingly branched, mostly 40–60  $\mu\text{m}$  in diameter, composed of segments 1.0–1.5 times as long as the diameter, arising at irregular intervals in a strictly endogenous manner; lateral branches exogenous or endogenous, independent of trichoblasts, which they replace; pericentral cells 4, ecorticate; trichoblasts mostly infrequent to wanting, with 1 or 2 dichotomies, 250–480  $\mu\text{m}$  long, soon deciduous.

Tetrasporangia 50–60  $\mu\text{m}$  in diameter, in straight series, somewhat distending the segments; cystocarps ovoid, 150–190  $\mu\text{m}$  in diameter; spermatangial stichidia cylindrical, without sterile apices, arising from an entire trichoblast primordium.

TYPE-LOCALITY.—“Ad littus americanae tropica,” probably on the Pacific coast of Mexico.

HOLOTYPE.—A collection by Liebmann in the Agardh Herbarium (LD).

GULF OF CALIFORNIA DISTRIBUTION.—Intertidal rocks, Puerto Peñasco; Isla Angel de la Guarda; Isla Partida; Isla Turner; Isla Estanque.

PACIFIC COAST DISTRIBUTION.—British Columbia, Canada, to Costa Rica. Widely distributed in the northern Pacific and western Atlantic oceans.

SPECIMEN STUDIED.—SONORA: JN-5098a, collected with *P. simplex* (JN-5098b), midtidal rocks, Playa Estación, Puerto Peñasco, 24 Mar 1974.

REMARKS.—This species is found growing alone or with other algae, often forming a continuous turf on intertidal rocks. Dawson (1944:332; 1954b:161) reported this taxon from three islands of Las Islas de la Cintura under the name of *Lophosiphonia villum* (J. G. Agardh) Setchell & Gardner. Our new collection, moves the known distribution northward to Puerto Peñasco from Puerto Refugio, Isla Angel de la Guarda (Dawson, 1944:333, 1963:422).

The reasons for placing this species in *Polysiphonia* rather than in *Lophosiphonia* are given by Hollenberg (1968a:56). In this connection, the criteria listed by Hollenberg indicate that *Lophosiphonia mexicana* described by Dawson (1944:333) cannot be properly assigned to that genus. Further collections of “*L. mexicana*” are needed before a definitive disposition of the taxon can be made.

*Polysiphonia simplex* Hollenberg

FIGURE 9

*Polysiphonia simplex* Hollenberg, 1942b:782, fig. 18; 1961:364, pl. 5: fig. 1.—Dawson, 1944:331; 1951:53; 1954a:6; 1954b:160; 1966:29.—Abbott and Hollenberg, 1976:694, fig. 641.

*Polysiphonia ferulacea* sensu Segi, 1951:209 [in part, with reference to Pacific North American specimens cited by Hollenberg, 1942b:782 only; not *Polysiphonia ferulacea* Suhr in J. G. Agardh, 1863:980].

DESCRIPTION.—Thalli medium to dark brownish red, drying nearly black; usually saxicolous, commonly forming dense mats often of considerable extent, alone or mixed with other algae, or occa-




FIGURE 9.—*Polysiphonia simplex*: A, upper portion of the thallus showing the delicate trichoblasts (JN-5098B); B, upper portion of a cystocarpic thallus (JN-5038).

sionally epiphytic; with creeping basal branches 250–360  $\mu\text{m}$  in diameter, composed of short segments, and attached by numerous unicellular rhizoids, which often have digitate apieces and are always cut off as separate cells from the proximal end of the pericentral cells; erect branches 1–3(–7) cm high and 160–250  $\mu\text{m}$  in diameter near the base, arising in an exogenous assurgent manner; main axes usually distinct but not prominent, sparingly branched; branches exogenous, radially directed, of several orders; trichoblasts delicate, to 730  $\mu\text{m}$  long with 1 or 2 dichotomies, 1 per segment in spiral sequence with  $\frac{1}{4}$  divergence, soon deciduous, leaving persistent scar-cells; branches replacing trichoblasts in the spiral (arising from an entire trichoblast primordium).

Tetrasporangia up to 70  $\mu\text{m}$  in diameter, in spiral series in the ultimate branches, more or less distending the segments; cystocarps ovoid-globular, 300–

350  $\mu\text{m}$  in diameter, spermatangial stichidia 100–170  $\mu\text{m}$  long, 35–40  $\mu\text{m}$  in diameter, comprising one primary branch of a trichoblast; without sterile apex.

TYPE-LOCALITY.—On rocks, Laguna Beach, Orange County, California.

HOLOTYPE.—H 2115, 14 May 1937, in herbarium of G. J. Hollenberg (now at US; isotype at GMS).

GULF OF CALIFORNIA DISTRIBUTION.—Intertidal, growing alone or with other turf algae forming mats; Puerto Peñasco to Cabeza Ballena.

PACIFIC COAST DISTRIBUTION.—Southern California to southern Mexico. Costa Rica.

SPECIMENS STUDIED.—SONORA: D 27173, Bahía Cholla, mud-flat, 5 Apr 1966; H 67-64, tetrasporangial, cystocarpic, Playa Tucson, Punta Pelicano, 9 Apr 1967; JN-5038, cystocarpic, low intertidal, Punta Pelicano, 17 Mar 1974; JN-5098b, mid-intertidal rock platform, Playa Estación, Puerto Peñasco, legit J. N. and K. B., 24 Mar 1974.

REMARKS.—This species is reported by Dawson (1944:331) and Hollenberg (1961:364) from a number of Gulf localities. With the recent collections from Puerto Peñasco, we now know *P. simplex* to be widely distributed throughout the Gulf.

*Polysiphonia sonorensis* Hollenberg

FIGURE 2C

*Polysiphonia sonorensis* Hollenberg, 1942b:779, figs. 5, 20.—Dawson, 1944:330; 1954b:161.—Hollenberg, 1961:365, pl. 5: fig. 2.

DESCRIPTION.—Densely tufted algae of a terracotta color, not strongly adhering to paper on drying, 4–5 cm high from prostrate creeping branches 120–175  $\mu\text{m}$  in diameter attached by unicellular rhizoids, which arise as outgrowths of the center of the pericentral cells, from which they are not cut off as separate cells; erect branches arising mostly endogenously or sometimes exogenously from the prostrate branches, 70–100  $\mu\text{m}$  in diameter, composed of segments mostly 1.5–2.0 times as long as the diameter; branching dense, the branches arising exogenously at irregular intervals of mostly 1–20 segments and mostly at an acute angle with the parent branch; main axes indistinct; branches replacing trichoblasts; trichoblasts numerous, delicate, to 1 mm long, arising at irregular intervals, with 1 or 2 dichotomies, soon deciduous, leaving inconspicuous scar-cells.

Reproductive material has not been found.

TYPE-LOCALITY.—Afloat along the southeastern shore of Bahía Empalme, Sonora, Gulf of California.

HOLOTYPE.—Francis Drouet and Donald Richards 3426a (F); isotype: herbarium of G. J. Hollenberg (now at GMS).

GULF OF CALIFORNIA DISTRIBUTION.—Intertidal; Bahía Empalme to Guaymas.

SPECIMENS STUDIED.—SONORA: in addition to the type, there is one other collection (Francis Drouet and Donald Richards-3144) from rocks between tide marks on NW shore of harbor at Guaymas.

REMARKS.—KNOWN only from two specimens, this species is apparently endemic to the Gulf. It was not found in the recent collections.

*Polysiphonia sphaerocarpa* var. *cheloniae*,  
new variety

FIGURES 4A–C, 6A

DESCRIPTION.—Plantae ad 3.5 cm altae, a crista basali rhizoideorum per rhizoidea e ramis assurgentibus suppletorum, ortae; rhizoidea per muros curvatos a finibus proximalibus cellulorum pericentralium ut cellulae discretae abscissa; cellulae pericentrales 4; ramificatio pseudodichotoma; rami principales ad 300  $\mu\text{m}$  crassi, e segmentis 1.0–2.5 (–3.0) plo longioribus quam lata compositi; segmentis in ramis iunioribus turgidulis; trichoblastae ad 400  $\mu\text{m}$  longae, una in unoquoque segmento, 1–3 dichotomias habentes, ad apices delicatos attenuatae; rami e primordio integro trichoblastae intervalis fere 5–12 segmentorum enascentes, basim versus angusti.

Tetrasporangia ignota; cystocarpi globosi, ad 370  $\mu\text{m}$  diametro; stichidia spermatangialia 160–175  $\mu\text{m}$  longa, 40–48  $\mu\text{m}$  diametro, cylindrica aut fusiformia, fere sine cacumine sterili, ut furca primaria trichoblastae enascentia.

Thalli to 3.5 cm high, from a basal tuft of rhizoids supplemented by rhizoids from assurgent branches; rhizoids cut off as separate cells from the proximal end of the pericentral cells; pericentral cells 4, ecorticate; branching pseudodichotomous; main branches to 300  $\mu\text{m}$  in diameter, composed of segments 1.0–2.5 (–3.0) times as long as the diameter; branches arising from an entire trichoblast primordium in intervals of mostly 5–12 segments, narrowed toward the base; segments in younger branches turgid; trichoblasts one per segment to 400  $\mu\text{m}$  long, with 1–3 dichotomies, tapering to delicate apices.

Tetrasporangia unknown; cystocarps globular, to 370  $\mu\text{m}$  in diameter; spermatangial stichidia 160–175  $\mu\text{m}$  long, 40–48  $\mu\text{m}$  in diameter, cylindrical to fusiform, mostly without sterile apices at maturity, arising as a primary fork of a trichoblast.

TYPE-LOCALITY.—On green sea turtle, *Chelonia mydas* L., in Canal de Infiernillo between Campo Ona and Campo Viboras, Sonora, opposite the east coast of Isla Tiburón. Gulf of California.

HOLOTYPE.—J. Norris-4801, cystocarpic (US).

GULF OF CALIFORNIA DISTRIBUTION.—Canal de Infiernillo, between Sonora and Isla Tiburón.




FIGURE 10.—*Polysiphonia* species: A, upper portion of a spermatangial thallus (JN-2998); B, branch tip, showing the spermatangial stichidia, which originates from an entire trichoblast primordium (JN-2998); C, upper portion, showing the usual absence of trichoblasts (JN-3265); D, upper portion of a tetrasporangial thallus, showing the developing tetrasporangia initials in the characteristic spiral sequence (JN-3265).

**SPECIMENS STUDIED.**—SONORA: all collections of this alga were made 21 Jan 1974 in the Infiernillo Channel between Campo Ona and Campo Viboras, Sonora (opposite the east coast of Isla Tiburón). They were collected by R. S. Felger and E. W. Moser from medium size to young adult green sea turtles (*Chelonia mydas*), which were harpooned by Guadalupe Lopez, a Seri Indian. Five collections were made: JN-4801, cystocarpic, holotype (US) was taken from the carapace of female turtle number 4, harpooned at approximately 9 m depth; JN-4765, partly spermatangial, was from the dorsal surface of the head and flippers of another female turtle number 2, harpooned at approximately 7.6 m depth; JN-4773, sterile, was collected from the carapace of female turtle number 1, taken from approximately 9 m depth; JN-4777, immature cystocarpic, was from the carapace of turtle number 1; and JN-4794, sterile, was from the dorsal surface of the head and front flippers of female turtle number 1.

**REMARKS.**—This alga seems very close in its general features to *P. sphaerocarpa* Boergesen (1918: 271) and especially to var. *sphaerocarpa* as described by Hollenberg (1968a:90) from the central and western Pacific Ocean. It differs from that variety in the substratum (i.e., the green sea turtle), and the consequent habitat. It is also somewhat taller than previously described specimens of that taxon.

At least two other species of *Polysiphonia* have been described as growing on sea turtles. *Polysiphonia tsudana*, collected from a species of sea turtle (*Chelonia*) taken near Laysan Island, Hawaii, was described by Hollenberg (1968b:205). It is smaller than the Gulf of California taxon, with branches much smaller in diameter and having shorter segments. Furthermore, the Laysan specimens of *P. tsudana* had a variable number of pericentral cells and the spermatangial stichidia arise from the entire trichoblast primordium, rather than from a primary branch as in *P. sphaerocarpa* var. *cheloniae* from the northern Gulf. *Polysiphonia carettia*, also described by Hollenberg (1971:15), was found growing on the carapace of a Loggerhead sea turtle (*Caretta caretta* L.), taken at the west end of Santa Catalina Island, off the coast of southern California. This alga is likewise smaller than the northern Gulf of California specimens. It differs, furthermore, in that the branches of the

California *P. carettia* are all or mostly all cicatrigenous in origin, arising from the scar-cells. Finally the rhizoids are cut off as separate cells in the Gulf of California specimens, but remain in open connection with the pericentral cells in the case of the California species.

### *Polysiphonia* species

FIGURE 10

**DESCRIPTION.**—Thalli to 3 cm high, repeatedly pseudodichotomously branched at irregular intervals of mostly 8–10 segments; main axes to 320  $\mu$ m in diameter below, composed of segments 1.0–1.5 times as long as the diameter; pericentral cells 4, ecorticate; rhizoids not observed; trichoblasts very rudimentary or absent, scar-cells 1 per segment in spiral sequence.

Tetrasporangia in spiral series in the smaller branches; cystocarps not seen; spermatangial stichidia arising from an entire trichoblast primordium.

**SPECIMENS STUDIED.**—BAJA CALIFORNIA DEL NORTE: two collections are assigned to this entity: JN-2998b, spermatangial, from 7.6 m depth, Isla la Ventana, Bahía de Los Angeles, 21 May 1972; and JN-3265b, tetrasporangial, growing on the "pen shell" (*Pinna rugosa*), 3–5 m depth, Bahía San Francisco, 24 May 1972.

**REMARKS.**—The manner of origin of spermatangial stichidia is very much like that in *P. pacifica* and *P. scopulorum*, but the presence of a scar-cell or a trichoblast on each segment is not a feature normally associated with the origin of spermatangial stichidia from an entire trichoblast primordium as in the last named species. Also there are differences in the arrangement of the tetrasporangia. In *P. pacifica* and *P. scopulorum* they are in a straight series, whereas in our Gulf of California collections they are arranged in a spiral series (Figure 10D). It seems probable that these specimens represent an undescribed species. Further collections are needed to determine their taxonomic status.



## Literature Cited

- Abbott, I. A., and G. J. Hollenberg  
 1976. *Marine Algae of California*. xii + [2] + 827 pages. Stanford: Stanford University Press.
- Agardh, J. G.  
 1863. *Species, genera et ordines Algarum, seu descriptiones succinctae specierum generum et ordinum, quibus algarum regnum constituitur*. Volume 2, part 3, pages 701–1291. Lund: G. W. K. Gleerup.
- Boergesen, F.  
 1918. The Marine Algae of the Danish West Indies, III: Rhodophyceae. *Danske Botanisk Arkiv*, 3:241–304.
- Dawson, E. Y.  
 1944. The Marine Algae of the Gulf of California. *Allan Hancock Pacific Expeditions*, 3(10):189–454.  
 1949. Resultados preliminares de un reconocimiento de las algas marinas de la costa Pacifica de México. *Revista de la Sociedad Mexicana de Historia Natural*, 9(3/4): 215–254.  
 1950. A Note on the Vegetation of a New Coastal Upwelling Area of Baja California. *Journal of Marine Research*, 9(2):66–68.  
 1951. A Further Study of Upwelling and Associated Vegetation along Pacific Baja California, Mexico. *Journal of Marine Research*, 10:39–58.  
 1954a. The Marine Flora of Isla San Benedicto Following the Volcanic Eruption of 1952–1953. *Allan Hancock Foundation Publications Occasional Paper*, 16:1–13.  
 1954b. Resúmen de las investigaciones recientes sobre algas marinas de la costa Pacifica de Mexico, con una sinopsis de la literatura, sinonimia y distribución de las especies descritas. *Revista de la Sociedad Mexicana de Historia Natural*, 13:97–197 + i-x [reprint of 1953 with corrections, index, pagination, and addenda].  
 1957. Notes on Eastern Pacific Insular Marine Algae. *Los Angeles County Museum Contributions in Science*, 8:1–8.  
 1959. Marine Algae from the 1958 Cruise of the *Stella Polaris* in the Gulf of California. *Los Angeles County Museum Contributions in Science*, 27:1–39.  
 1963. Marine Red Algae of Pacific Mexico, Part 8, Ceramiales: Dasyaceae, Rhodomelaceae. *Nova Hedwigia*, 6:401–481.  
 1966. Marine Algae in the Vicinity of Puerto Peñasco, Sonora, Mexico. *Gulf of California Field Guide Series*, 1:iii + 57. Tucson: University of Arizona.
- Dawson, E. Y., C. Acleto O., and N. Foldvik  
 1964. The Seaweeds of Peru. *Beihefte zur Nova-Hedwigia*, 13: [5] + 111 pages.
- Dawson, E. Y., M. Neushul, and R. D. Wildman  
 1960a. Seaweeds Associated with Kelp Beds along Southern California and Northwestern Mexico. *Pacific Naturalist*, 1(14):1–81.  
 1960b. New Records of Sublittoral Marine Plants from Pacific Baja California. *Pacific Naturalist*, 1(19):1–30.
- Felger, R. S., K. Clifton, and P. J. Regal  
 1976. Winter Dormancy in Sea Turtles: Independent Discovery and Exploitation in the Gulf of California by Two Local Cultures. *Science*, 191:283–285.
- Gardner, N. L.  
 1927. New Rhodophyceae from the Pacific Coast of North America, VI. *University of California Publications, Botany*, 14:99–138.
- Greville, R. K.  
 1824. *Flora Edinensis*. lxxxii + 478 pages. Edinburgh: W. Blackwood.
- Hariot, P.  
 1891. Liste des algues marines reportées de Yokosuka (Japon) par M. le Dr. Savatier. *Memoires de la Société des Sciences Naturelles et Mathématiques de Cherbourg*, 27:211–230.
- Harvey, W. H.  
 1847. *Nereis Australis, or Algae of the Southern Ocean*. [2] + viii + 124 pages. London: Reeve Brothers.  
 1853. *Nereis Boreali-Americana; or, Contributions towards a History of the Marine Algae of the Atlantic and Pacific Coasts of North America, Part II: Rhodospirameae*. 258 pages. Washington, D. C.: Smithsonian Institution.
- Hollenberg, G. J.  
 1942a. Phycological Notes, I. *Bulletin of the Torrey Botanical Club*, 69:528–538.  
 1942b. An Account of the Species of *Polysiphonia* on the Pacific Coast of North America, I: Oligosiphonia. *American Journal of Botany*, 29:772–785.  
 1944. An Account of the Species of *Polysiphonia* on the Pacific Coast of North America, II: Polysiphonia. *American Journal of Botany*, 31:474–483.  
 1948. Notes on Pacific Coast Marine Algae. *Madroño*, 1948:155–162.  
 1961. Marine Red Algae of Pacific Mexico, Part 5: The Genus *Polysiphonia*. *Pacific Naturalist*, 2(6):345–375.  
 1968a. An Account of the Species of *Polysiphonia* of the Central and Western Tropical Pacific Ocean, I: Oligosiphonia. *Pacific Science*, 22(1):56–98.  
 1968b. An Account of the Species of the Red Alga *Polysiphonia* of the Central and Western Tropical Pacific Ocean, II: Polysiphonia. *Pacific Science*, 22(2):198–207.  
 1971. Phycological Notes, V: New Species of Marine Algae from California. *Phycologia*, 10(1):11–16.

- Holmgren, P. K., and W. Keuken  
 1974. *Index Herbariorum, Part I: The Herbaria of the World*. Sixth edition, vii + 397 pages. Utrecht, Netherlands: Oosthoek, Scheltema & Holkema.
- Hooker, W. J.  
 1833. Mosses, Hepaticae, Lichens, Characeae and Algae. Part I of volume 5 (*Class XXIV, Cryptogamia*) in *The English Flora of Sir James Edward Smith*. x + 4 + 432 pages. London: Longman, Rees, Orme, Brown, Green & Longman. [Listed under "J. E. Smith" in most libraries.]
- Howe, M. A.  
 1914. The Marine Algae of Peru. *Memoirs of the Torrey Botanical Club*, 15:1-185.
- Kuetzing, F. T.  
 1863. *Tabulae Phycologicae*. Volume 13, [i] + 31 pages. Nordhausen.  
 1864. *Tabulae Phycologicae*. Volume 14, [i] + 35 pages. Nordhausen.
- Kylin, H.  
 1941. Californische Rhodophyceen. *Lunds Universitets Arsskrift, N. F. Avd. 2*, 37(2):1-51; and *Kunglig Fysiografiska Sällskapets Handlingar*, new series, 52(2):1-51.
- Montagne, C.  
 1842. Troisième centurie de plantes cellulaires exotiques nouvelle, part 2. *Annales des Sciences Naturelles, Botanique*, series 2, 18:241-282.
- Norris, J. N.  
 1972. Marine Algae from the 1969 Cruise of "Makrele" to the Northern Part of the Gulf of California. *Boletín de la Sociedad Botánica de México*, 32:1-30.
- Norris, J. N., and K. E. Bucher  
 1976. New Records of Marine Algae from the 1974 R/V *Dolphin* Cruise to the Gulf of California. *Smithsonian Contributions to Botany*, 34:1-22.
- Reinsch, P. F.  
 1888. Species et genera nova Algarum ex insula Georgia Australi. *Berichte der Deutschen Botanischen Gesellschaft*, 6:144-156.
- Segi, T.  
 1951. Systematic Study of the Genus *Polysiphonia* from Japan and Its Vicinity. *Journal of the Faculty of Fisheries, Prefectural University of Mie*, 1(2):169-272.
- Setchell, W. A., and N. L. Gardner  
 1903. Algae of Northwestern America. *University of California Publications, Botany*, 1:165-418.  
 1924. The Marine Algae: Expedition of the California Academy of Sciences to the Gulf of California in 1921. *Proceedings of the California Academy of Sciences*, fourth series, 12:695-949.  
 1930. Marine Algae of the Revillagigedo Islands Expedition in 1925. *Proceedings of the California Academy of Sciences*, fourth series, 19:109-215.
- Smith, G. M.  
 1944. *Marine Algae of the Monterey Peninsula, California*. ix + 622 pages. Stanford: Stanford University Press.  
 1969. *Marine Algae of the Monterey Peninsula, California*. Second edition (incorporating the 1966 *Supplement* by G. J. Hollenberg and I. A. Abbott), x + 752 pages. Stanford: Stanford University Press.
- Taylor, W. R.  
 1945. Pacific Marine Algae of the Allan Hancock Expeditions to the Galápagos Islands. *Allan Hancock Pacific Expeditions*, 12: iv + 528 pages.
- Womersley, H. B. S.  
 1950. The Marine Algae of Kangaroo Island, III: List of Species, I. *Transactions of the Royal Society of Southern Australia*, 73(2):137-197.

# Index

(Synonyms and principle entries in *italics*; new variety in SMALL CAPITALS)

Lophosiphonia, 14  
  *mexicana*, 14  
  *scopulorum*, 14  
  *villum*, 14

Polysiphonia, 2 (key)  
  *acuminata*, 4  
  *californica*, 12  
  *concinna*, 7  
  *confusa*, 2, 3 (fig.)  
  *eastwoodae*, 9  
  *ferulacea*, 14  
  *flaccidissima*, 4, 5 (fig.)  
  *inconspicua*, 2  
  *johnstonii* var. *concinna*, 5 (fig.), 7, 8 (fig.)  
  *johnstonii* var. *johnstonii*, 4, 5 (fig.), 6 (fig.)  
  *marchantae*, 4  
  *masonii*, 5 (fig.), 8

*minutissima*, 12  
  *mollis*, 9, 10 (fig.)  
  *pacifica* var. *delicatula*, 10, 11 (fig.)  
  *paniculata*, 10 (fig.), 11, 12 (fig.)  
  *richardsoni*, 4  
  *savatieri*, 12, 13 (fig.)  
  *scopulorum* var. *villum*, 13 (fig.), 14  
  *simplex*, 14, 15 (fig.)  
  *sinicola*, 4  
  *snyderae*, 9  
  *snyderae* var. *heteromorpha*, 9  
  *snyderae* var. *intricata*, 9  
  *sonorensis*, 5 (fig.), 16  
  *species*, 17, 18 (fig.)  
  *sphaerocarpa* var. *CHELONIAE*, 8 (fig.), 11 (fig.), 16  
  *tongatensis*, 9  
  *tongatensis* var. ? , 9  
  *villum*, 14







## REQUIREMENTS FOR SMITHSONIAN SERIES PUBLICATION

**Manuscripts** intended for series publication receive substantive review within their originating Smithsonian museums or offices and are submitted to the Smithsonian Institution Press with approval of the appropriate museum authority on Form SI-36. Requests for special treatment—use of color, foldouts, casebound covers, etc.—require, on the same form, the added approval of designated committees or museum directors.

**Review** of manuscripts and art by the Press for requirements of series format and style, completeness and clarity of copy, and arrangement of all material, as outlined below, will govern, within the judgment of the Press, acceptance or rejection of the manuscripts and art.

**Copy** must be typewritten, double-spaced, on one side of standard white bond paper, with 1 $\frac{1}{4}$ " margins, submitted as ribbon copy (not carbon or xerox), in loose sheets (not stapled or bound), and accompanied by original art. Minimum acceptable length is 30 pages.

**Front matter** (preceding the text) should include: **title page** with only title and author and no other information, **abstract page** with author/title/series/etc., following the established format, **table of contents** with indents reflecting the heads and structure of the paper.

**First page of text** should carry the title and author at the top of the page and an unnumbered footnote at the bottom consisting of author's name and professional mailing address.

**Center heads** of whatever level should be typed with initial caps of major words, with extra space above and below the head, but with no other preparation (such as all caps or underline). Run-in paragraph heads should use period/dashes or colons as necessary.

**Tabulations** within text (lists of data, often in parallel columns) can be typed on the text page where they occur, but they should not contain rules or formal, numbered table heads.

**Formal tables** (numbered, with table heads, boxheads, stubs, rules) should be submitted as camera copy, but the author must contact the series section of the Press for editorial attention and preparation assistance before final typing of this matter.

**Taxonomic keys** in natural history papers should use the aligned-couplet form in the zoology and paleobiology series and the multi-level indent form in the botany series. If cross-referencing is required between key and text, do not include page references within the key, but number the keyed-out taxa with their corresponding heads in the text.

**Synonymy** in the zoology and paleobiology series must use the short form (taxon, author, year:page), with a full reference at the end of the paper under "Literature Cited." For the botany series, the long form (taxon, author, abbreviated journal or book title, volume, page, year, with no reference in the "Literature Cited") is optional.

**Footnotes**, when few in number, whether annotative or bibliographic, should be typed at the bottom of the text page on which the reference occurs. Extensive notes must appear at the end of the text in a notes section. If bibliographic footnotes are required, use the short form (author/brief title/page) with the full reference in the bibliography.

**Text-reference system** (author/year/page within the text, with the full reference in a "Literature Cited" at the end of the text) must be used in place of bibliographic footnotes in all scientific series and is strongly recommended in the history and technology series: "(Jones, 1910:122)" or ". . . Jones (1910:122)."

**Bibliography**, depending upon use, is termed "References," "Selected References," or "Literature Cited." Spell out book, journal, and article titles, using initial caps in all major words. For capitalization of titles in foreign languages, follow the national practice of each language. Underline (for italics) book and journal titles. Use the colon-parentheses system for volume/number/page citations: "10(2):5-9." For alignment and arrangement of elements, follow the format of the series for which the manuscript is intended.

**Legends** for illustrations must not be attached to the art nor included within the text but must be submitted at the end of the manuscript—with as many legends typed, double-spaced, to a page as convenient.

**Illustrations** must not be included within the manuscript but must be submitted separately as original art (not copies). All illustrations (photographs, line drawings, maps, etc.) can be intermixed throughout the printed text. They should be termed **Figures** and should be numbered consecutively. If several "figures" are treated as components of a single larger figure, they should be designated by lowercase italic letters (underlined in copy) on the illustration, in the legend, and in text references: "Figure 9**b**." If illustrations are intended to be printed separately on coated stock following the text, they should be termed **Plates** and any components should be lettered as in figures: "Plate 9**b**." Keys to any symbols within an illustration should appear on the art and not in the legend.

**A few points of style:** (1) Do not use periods after such abbreviations as "mm, ft, yds, USNM, NNE, AM, BC." (2) Use hyphens in spelled-out fractions: "two-thirds." (3) Spell out numbers "one" through "nine" in expository text, but use numerals in all other cases if possible. (4) Use the metric system of measurement, where possible, instead of the English system. (5) Use the decimal system, where possible, in place of fractions. (6) Use day/month/year sequence for dates: "9 April 1976." (7) For months in tabular listings or data sections, use three-letter abbreviations with no periods: "Jan, Mar, Jun," etc.

**Arrange and paginate sequentially EVERY sheet of manuscript**—including ALL front matter and ALL legends, etc., at the back of the text—in the following order: (1) title page, (2) abstract, (3) table of contents, (4) foreword and/or preface, (5) text, (6) appendixes, (7) notes, (8) glossary, (9) bibliography, (10) index, (11) legends.

SMITHSONIAN INSTITUTION LIBRARIES



3 9088 01004 5425

